



## **Curtis J Kephart, MD**

**Orthopedic Surgery and Sports Medicine**  
**FL License: 116858**

# Proximal Humerus Fracture Rehabilitation Protocol

### **Unstable Proximal Humerus Fracture**

- This protocol is for patients with unstable proximal humerus fractures.

**Goal:** Maintain ROM without displacing the proximal humerus fracture.

- Due to variable stability of the fracture, check for specifications and precautions ordered by the orthopedist.

### **Phase I (0 to 21 days)**

- Begin elbow, wrist and hand active ROM.
- A sling and swath, or Velpeau with an axillary pad should be used when not doing physical therapy.
- After 14 days, begin pendulum exercises (clockwise and counterclockwise).
- Establish a home exercise program so patient is performing exercises 3-5 times per day for 10 minutes each session.

### **Phase II (3 to 6 weeks)**

- Begin supine ER with a cane. Fifteen to 20 degrees of abduction are permitted if the patient is more comfortable.
- Begin active assisted forward elevation (FE).
- Perform pulley exercises and teach for home program.
- Perform isometric exercises for IR, ER, extension, and abduction.

### **Phase III (7 weeks to 2 months)**

- Begin supine active FE. Progressively increase patients position from supine to erect during FE exercises.
- Use Therabands of progressive strengths for IR, ER, anterior, middle and posterior deltoid.
- Begin flexibility and stretching exercises to progressively increase ROM in all positions (i.e. towel behind back, finger walking up the wall, etc.).

## **Curtis J Kephart, MD**

**Orthopaedic Surgery and Sports Medicine**

13590 Jog Road Suite 7  
Delray Beach, FL 33446  
(561) 637-4200 OFFICE  
(561)-637-3222 FAX

9970 Central Park Blvd Suite 300  
Boca Raton, FL 33428  
(561)488-2200 OFFICE  
(561)488-1064 FAX