

Curtis J Kephart, MD Orthopedic Surgery and Sports Medicine (561) 637-4200 OFFICE (561) 637-3222 FAX FL License: 116858

Patient Sticker Here

Date

<u>PHYSICAL THERAPY PRESCRIPTION:</u> Open Reduction Internal Fixation of the Ankle Rehabilitation Protocol for Physical Therapy

Week 0-2:

Patient non weightbearing in splint with ankle in equinus At 2 weeks, splint removed, wound checked, and placed into CAM boot

Weeks 2-4:

Skin staples/sutures removed Continue non weightbearing in boot When incisions healed, begin passive / active dorsiflexion and plantar flexion stretch

Week 4-8:

PWB in boot from 4-6 weeks. Can WBAT in boot after 6 weeks. Boot discontinued at 8 weeks Passive / active dorsiflexion and plantar flexion stretch Inversion / Eversion ROM exercises Begin dorsiflexion and plantar flexion isometrics and progress to isotonics Can start riding a stationary bicycle Should be walking on a treadmill with wean up to 3.5 mph

Week 9-11: Advance strengthening

Week 12 +: Can begin jogging, stairmaster Add isokinetics Increase strengthening, endurance, proprioception, flexibility exercise Initiate sport specific drills with gradual return to athletics at or after 16 weeks

Please send progress notes.

Physician's Signature:_____

Curtis J Kephart, MD